

$X(3900)^{\pm}$ $I(J^P) = ?(1^+)$

Seen as a peak in the invariant mass distribution of the $J/\psi\pi^{\pm}$ system by BES3 (ABLIKIM 13T) in $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ at c.m. energy of 4.42 GeV and by radiative return from e^+e^- collisions at \sqrt{s} from 9.46 to 10.86 GeV at BELLE (LIU 13B). Angular analysis of ABLIKIM 14A favors the $J^P = 1^+$ assignment. Needs confirmation.

NODE=M210

 $X(3900)^{\pm}$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3888.7±3.4 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
3883.9±1.5±4.2	1.2k	1,2 ABLIKIM	14A BES3	$e^+e^- \rightarrow \pi^{\pm}(D\bar{D}^*)^{\mp}$
3899.0±3.6±4.9	307 ± 48	1 ABLIKIM	13T BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
3894.5±6.6±4.5	159 ± 49	1 LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
3886 ± 4 ± 2	81 ± 16	3 XIAO	13A	4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ Neglecting interference between the $X(3900)$ and non-resonant continuum.² With estimated statistical significance of more than 18σ .³ For $M^2(\pi^+\pi^-) < 0.65$ GeV 2 . Signal has 5.7σ significance including systematic uncertainties. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.


NODE=M210M

NODE=M210

NODE=M210M;LINKAGE=A

NODE=M210M;LINKAGE=C

NODE=M210M;LINKAGE=B



NODE=M210W

NODE=M210W

NODE=M210W;LINKAGE=A

NODE=M210W;LINKAGE=C

NODE=M210W;LINKAGE=B

 $X(3900)^{\pm}$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
35 ± 7 OUR AVERAGE				
24.8± 3.3±11.0	1.2k	1,2 ABLIKIM	14A BES3	$e^+e^- \rightarrow \pi^{\pm}(D\bar{D}^*)^{\mp}$
46 ± 10 ± 20	307 ± 48	1 ABLIKIM	13T BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
63 ± 24 ± 26	159 ± 49	1 LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
37 ± 4 ± 8	81 ± 16	3 XIAO	13A	4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ Neglecting interference between the $X(3900)$ and non-resonant continuum.² With estimated statistical significance of more than 18σ .³ For $M^2(\pi^+\pi^-) < 0.65$ GeV 2 . Signal has 5.7σ significance including systematic uncertainties. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

X(3900) $^\pm$ DECAY MODES

NODE=M210215;NODE=M210

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\pi^\pm$	seen
Γ_2 $h_c\pi^\pm$	not seen
Γ_3 $(D\bar{D}^*)^\pm$	seen

X(3900) $^\pm$ BRANCHING RATIOS

$\Gamma(J/\psi\pi^\pm)/\Gamma_{\text{total}}$	Γ_1/Γ			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	307 ± 48	ABLIKIM	13T BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(h_c\pi^\pm)/\Gamma_{\text{total}}$	Γ_2/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	ABLIKIM	13X BES3	$e^+e^- \rightarrow h_c\pi^\pm$

$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi\pi^\pm)$	Γ_3/Γ_1			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.2\pm1.1\pm2.7	1.2k	¹ ABLIKIM	14A BES3	$e^+e^- \rightarrow \pi^\pm (D\bar{D}^*)^\pm$

¹ Assuming the same origin of the $(D\bar{D}^*)^\pm$ and $\pi^\pm J/\psi$ decay modes.

X(3900)$^\pm$ REFERENCES				
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BES III Collab.) JP
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BES III Collab.)
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)

DESIG=1
DESIG=2
DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M210225

NODE=M210R01
NODE=M210R01

NODE=M210R02
NODE=M210R02

NODE=M210R03
NODE=M210R03

NODE=M210R03;LINKAGE=A

NODE=M210

REFID=55648
REFID=55409
REFID=55635
REFID=55410
REFID=55593